



Prof (Dr) Upendra Singh (M. Pharm, Ph.D) Om Group of Colleges, Roorkee, Haridwar (U.K.)

OFFICE: BUILDING No. 3/314, OFFICE-1, GAUSHALA ROAD, SHAMLI DISTRICT SHAMLI (U.P.) -

247776

Mobile: +91-9719638415

Email: arvindrkgit@gmail.com, www.phbeducation.com

D. Pharm Exit Exam - 2024

Course Name : D. Pharm

Year : Second Year

Subject Name: Pharmacology

Topic Name : Anti-depressant drugs

Anti-depressant drugs

MULTIPLE CHIOCE QUESTION

- **1.** Which of the following neurotransmitters is primarily targeted by most antidepressant drugs?
- A) Dopamine
- B) Serotonin
- C) Acetylcholine
- D) Glutamate

Answer: B) Serotonin

- **2.** Which class of antidepressants is known for inhibiting the reuptake of serotonin and norepinephrine?
- A) Selective serotonin reuptake inhibitors (SSRIs)
- B) Monoamine oxidase inhibitors (MAOIs)
- C) Tricyclic antidepressants (TCAs)
- D) Atypical antidepressants

Answer: A) Selective serotonin reuptake inhibitors (SSRIs)

- **3.** Which antidepressant class is associated with the inhibition of monoamine oxidase enzyme?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: C) MAOIs

- **4.** Which of the following is NOT a common side effect of SSRIs?
- A) Weight gain
- B) Sexual dysfunction
- C) Insomnia
- D) Nausea

Answer: A) Weight gain

- **5.** Bupropion is classified as:
- A) An SSRI
- B) A TCA
- C) A MAOI

D) An atypical antidepressant

Answer: D) An atypical antidepressant

- **6.** Which antidepressant class is associated with potentially dangerous interactions with certain foods rich in tyramine?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: C) MAOIs

- 7. Which neurotransmitter is primarily targeted by TCAs?
- A) Serotonin
- B) Dopamine
- C) Norepinephrine
- D) GABA

Answer: C) Norepinephrine

- **8.** Which antidepressant class is often considered the first-line treatment for depression due to their favorable side effect profile and safety?
- A) SSRIs
- B) MAOIs
- C) TCAs
- D) Atypical antidepressants

Answer: A) SSRIs

- **9.** Which of the following is a common side effect of TCAs?
- A) Weight loss
- B) Sedation
- C) Sexual dysfunction
- D) Hyponatremia

Answer: B) Sedation

- 10. Duloxetine belongs to which class of antidepressants?
- A) SSRIs
- B) TCAs
- C) SNRIs
- D) MAOIs

Answer: C) SNRIs (Serotonin-norepinephrine reuptake inhibitors)

- **11.** Which antidepressant is commonly used to help with smoking cessation? A) Venlafaxine
- B) Duloxetine
- C) Bupropion
- D) Mirtazapine

Answer: C) Bupropion

- **12.** Which class of antidepressants is known for causing significant anticholinergic side effects?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: B) TCAs

- **13.** Which antidepressant class has a potential risk of serotonin syndrome when combined with other serotonergic medications?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: A) SSRIs

- 14. Which of the following is NOT an atypical antidepressant?
- A) Mirtazapine
- B) Trazodone
- C) Vilazodone
- D) Fluoxetine

Answer: D) Fluoxetine

- **15.** Which antidepressant is often used for depression with insomnia due to its sedating properties?
- A) Fluoxetine
- B) Escitalopram
- C) Trazodone
- D) Sertraline

Answer: C) Trazodone

- **16.** Which of the following is a potential side effect of MAOIs when combined with certain foods or medications?
- A) Hypotension
- B) Hypertensive crisis
- C) Hyperglycemia
- D) Bradycardia

Answer: B) Hypertensive crisis

- **17.** Which antidepressant class is often considered safer for use in elderly patients due to their lower risk of adverse effects?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: A) SSRIs

- **18.** Which antidepressant is commonly used to treat depression in adolescents and young adults?
- A) Amitriptyline
- B) Paroxetine
- C) Imipramine
- D) Phenelzine

Answer: B) Paroxetine

- 19. Which of the following is NOT a common side effect of atypical antidepressants?
- A) Weight gain
- B) Dry mouth
- C) Sexual dysfunction
- D) Orthostatic hypotension

Answer: D) Orthostatic hypotension

- **20.** Which antidepressant class is associated with the highest risk of discontinuation syndrome upon abrupt cessation?
- A) SSRIs
- B) TCAs
- C) MAOIs
- D) Atypical antidepressants

Answer: A) SSRIs